

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

## Journal of Archaeological Science

journal homepage: <http://www.elsevier.com/locate/jas>

## Slaves as burial gifts in Viking Age Norway? Evidence from stable isotope and ancient DNA analyses

Elise Naumann <sup>a, \*</sup>, Maja Krzewińska <sup>b, c</sup>, Anders Götherström <sup>d</sup>, Gunilla Eriksson <sup>d</sup><sup>a</sup> Department of Archaeology, Conservation and History, University of Oslo, Oslo, Norway<sup>b</sup> Department of Biosciences, University of Oslo, Norway<sup>c</sup> Museum of Cultural History, University of Oslo, Norway<sup>d</sup> Archaeological Research Laboratory, Department of Archaeology and Classical Studies, Stockholm University, Sweden

## ARTICLE INFO

## Article history:

Received 29 May 2013

Received in revised form

24 August 2013

Accepted 28 August 2013

## Keywords:

Ancient DNA

Stable isotopes

Vikings

Slaves

Norway

## ABSTRACT

Ten Viking Age individuals from the northern Norwegian site at Flakstad were analysed for  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$  and ancient mitochondrial DNA fragments. The material derives from both single and multiple burials with individuals treated in different ways. The genetic analyses show that the individuals buried together were unlikely to be maternally related, and stable isotope analyses suggest different strata of society. It is, therefore, suggested that slaves may have been offered as grave gifts at Flakstad. A comparison with the remaining population from single graves shows that the presumed slaves had a diet similar to that of the common population, whereas the high status individuals in multiple graves had a diet different from both slaves and the common population. The results provide an insight into the subsistence of different social groups in a Viking Age society, exposing unexpected patterns of living conditions and food distribution.

© 2013 The Authors. Published by Elsevier Ltd. Open access under [CC BY-NC-SA license](http://creativecommons.org/licenses/by-nc-sa/4.0/).

## 1. Introduction

The Viking Age (A.D. 800–1030) was characterized by striking heterogeneity in burial customs, with regard to burial form, grave artefacts and treatment of the body. Among these, instances where two or more individuals were buried together stand out as a unique possibility to study social relations in the past. Graves with two or more individuals occur quite frequently throughout the Viking Age and all over the Viking World. The choice to bury persons together is hardly coincidental, but rather a deliberate action based on specific relationships between individuals. These relationships could be egalitarian, reflecting family members or persons otherwise closely connected. Alternatively the choice could involve human sacrifice or in other ways persons of different social rank, where one or more persons are intended to accompany the “main” burial. In Norway, few skeletons from inhumation burials remain due to poor preservation conditions, and thus indications of social relations, based on body treatment and artefact distribution, are often absent.

On the northern Norwegian Island of Flakstad a Late Iron Age (A.D. 550–1030) burial ground was located and partly excavated in the period 1980–1983. Human remains from ten individuals in three single burials, two double and one triple burial, were dated to the Viking Age ([The Schreiner Collection Database, 2013](#); [Sandmo, 1985a](#)). Although much of the contextual information has been lost due to farming activity, the burials showed unusual features. In each of the double/triple burials one individual was buried intact, while only post-cranial bones were found from the remaining individuals. The burial form at Flakstad may indicate graves containing master and slave, as crania from four individuals were absent from the graves. This interpretation is supported by a number of double burials found within Norse societies indicating slaves buried with their masters ([Andersen, 1960](#); [Hemmendorff, 1984](#); [Holck, 1997](#); [Holmquist Olausson, 1990](#); [Ramskou, 1965](#); [Skre, 1998](#); [Zachrisson, 2003](#)). The presumed presence of slaves is based on a number of factors such as maltreatment of the body, decapitation, binding of hands and feet and uneven distribution of grave gifts. At Flakstad the four accompanying individuals were seemingly decapitated, and thus the burials immediately appear to fit this pattern. However, as knowledge of the burial context from Flakstad is limited, the material calls for further investigation to study the social relations represented. Through application of stable isotope and ancient DNA (aDNA) analyses the presented

\* Corresponding author.

E-mail address: [elisena@gmail.com](mailto:elisena@gmail.com) (E. Naumann).

investigation aims at identifying dietary and genetic patterns that may highlight social relations reflected in differential treatment of the dead.

## 2. Material and methods

The burial site of Flakstad is situated on Flakstad Island, part of the Lofoten archipelago in northwestern Norway in Nordland County (Fig. 1). The burial site is situated close to a settlement area, which exhibits traces of contemporary agrarian activity. Parts of the burial site at Flakstad vicarage were excavated in the period 1980–1983 (Sandmo, 1985a,b). The ten individuals presented in this paper were all dated to the Viking Age based on burial artefacts. Graves 5858, 5860 and 5861 are single burials, and were excavated and documented by archaeologists. Burial artefacts consisted of weapons, comb, beads, garment accessories and animal bones, representing different levels of social wealth (Sandmo, 1985b). Graves 5863 and 5865 were registered as double burials (The Schreiner Collection Database, 2013). In grave 5864 a third individual was identified, but the remains are so fragmented that absence of the skull may be due to a general problem of preservation.

The skeletons in double/triple graves were buried in shallow graves close to the surface, and were partly disturbed by agricultural activity before archaeologists arrived. The disturbance was mainly due to the ongoing activity at the site, but earlier disturbance cannot be ruled out. Thus, the interpretation of multiple burials was questioned (Sandmo, 1985b). Information on

burial context is, therefore, poor and fragmented, and few burial artefacts were recovered from the double graves; two knives, a horse bit, a bead of amber, animal bones, parts of a whetstone and iron fragments. Unfortunately no information on artefact distribution within these burials exists. The skeletons were found in clusters, with an interval of approximately 30 m between each cluster. Each double/triple grave contained only one skull, but postcranial remains from two or three individuals. It seems unlikely that the missing skulls would be lost by coincidence in all four cases. In the osteological report no signs of decapitation were mentioned (Holck, 1983), although lack of evidence cannot serve as a strong argument. It should be noted that the heads might also have been removed post mortem. To ensure that cranial and postcranial material matched in the double/triple graves, samples from both long bones and mandible were taken in addition to teeth.

### 2.1. Stable isotopes

Food consumption is reflected in the isotopic composition of human tissue. Stable carbon and nitrogen isotope ratios are expressed as isotope values,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ , and reported relative to the standard PDB limestone for carbon and AIR atmospheric  $\text{N}_2$  for nitrogen. Carbon isotopes fractionate differently depending on the photosynthetic pathway, but also differ between marine and terrestrial environments (Chisholm et al., 1982). Since only the  $\text{C}_3$  pathway is relevant in an Arctic context,  $\delta^{13}\text{C}$  is here analysed to distinguish between terrestrial and marine food sources. Nitrogen




Fig. 1. Flakstad, situated in Nordland County.

isotopes fractionate for each step in the food chain (Minagawa and Wada, 1984), and the  $\delta^{15}\text{N}$  value therefore indicates trophic level. The  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in bone and dentine collagen mainly reflect the protein intake (Ambrose and Norr, 1993). Isotopic values in teeth reflect the period of tooth formation, and thus the childhood and adolescent years of the individual. Bone tissue, by contrast, remodels throughout life and therefore provides an average isotopic value reflecting the last 5–20 years, depending on various conditions, such as type of bone and age of the individual (Hedges et al., 2007; Sealy et al., 1995). Due to this variation, we have analysed both tooth dentine and bone to assess whether there has been continuity or changes in diet during the lifetime of an individual. Where possible, the cervix of the second molar, reflecting childhood (c. 7.5–10 years of age), was sampled for comparability.

Because of different ecological conditions such as salinity, temperature, distance from land and variations over time, isotopic values must always be considered in relation to the local fauna (Jones et al., 2012). In northern Europe it is often assumed that  $\delta^{13}\text{C}$  ratios range from approximately  $-22\text{‰}$  for a purely land based nutrition to  $-12\text{‰}$  for an exclusively marine diet (Chisholm et al., 1982). However,  $\delta^{13}\text{C}$  values from North Atlantic cod collagen emphasize the complexity of  $\delta^{13}\text{C}$  values in arctic Norway, as they range from  $-13\text{‰}$  to  $-17\text{‰}$  for migratory cod (Barrett et al., 2008). Rather than an absolute end-value for a purely marine protein intake, we therefore operate with a predicted  $\delta^{13}\text{C}$  range for humans consuming an exclusively marine diet in the northern regions of Norway. Available data documenting the  $\delta^{13}\text{C}$  offset between cod collagen (reference data) and muscle (consumed diet) is limited, though studies performed on modern gadid fish from Scotland showed on average a depletion of  $4\text{‰}$  in muscle tissue compared to collagen (Barrett et al., 2000a,b). Considering a trophic increase of  $5\text{‰}$  in  $\delta^{13}\text{C}$  values from diet to collagen in humans (Ambrose et al., 1997), we have taken into account a  $+1\text{‰}$  trophic-level effect for  $\delta^{13}\text{C}$  in the estimation of marine consumption.

The  $\delta^{15}\text{N}$  value increases by  $3\text{–}5\text{‰}$  for each trophic level in the food chain, and thus provides a distinction between herbivores and carnivores and between terrestrial and marine organisms, since the food chains are longer in marine environments (Bocherens and Drucker, 2003; Minagawa and Wada, 1984). The  $\delta^{15}\text{N}$  values of predatory marine mammals represent the highest levels in animals as they are at the top of the food chain. The values may thus range from  $6$  to  $8\text{‰}$  for a purely vegetarian diet to around  $20\text{‰}$  for a diet based exclusively on seal, as reflected in the analysis of a Greenland Inuit population (Schoeninger et al., 1983).  $\delta^{15}\text{N}$  values in a consumer's collagen reflect consumed diet plus trophic level, allowing direct interpretation of  $\delta^{15}\text{N}$ -results without offset correction (Ambrose et al., 1997).

Available fauna recovered from the Flakstad graves included specimens of cattle, sheep/goat, horse, dog and whale. All samples for stable isotopes were drilled with a dentist drill after removal of the surface to avoid contamination. Bone and dentine collagen was extracted according to Brown et al. (1988), which includes an ultrafiltration step to remove contaminants and degraded products of molecular sizes  $<30$  kDa. The extracted collagen was freeze-dried and the carbon and nitrogen isotope ratios determined using a Carlo Erba NC2500 analyser connected via a split interface to reduce the gas volume to a Finnigan MAT Delta + mass spectrometer, at the Stable Isotope Laboratory (SIL), Department of Geological Sciences, Stockholm University. The precision was  $\pm 0.15\text{‰}$  or better.

## 2.2. Ancient DNA

The ancient DNA analyses of the Flakstad human remains targeted mitochondrial DNA (mtDNA). The molecule has multiple

copies in human cells and is inherited from mother to child. The properties allow for tracing ancient human migrations and genetic affiliation through the maternal lineage (see Bandelt et al., 2006 for full overview). In order to investigate putative origin and possible maternal relationships among individuals from Flakstad, we have analysed bone samples from all 10 individuals presented here.

General guidelines for work with ancient DNA were followed at all times (Pääbo, 1989; Richards et al., 1995). Apart from an additional UV-radiation step, bone powder samples were prepared following Hagelberg and Clegg (1991). Between 20 and 50 mg of bone powder underwent DNA extraction with Qiagen DNeasy Blood & Tissue Kit (Qiagen) according to a modified manufacturer protocol. MtDNA was amplified between positions 16,050–16,392 in five overlapping fragments as described in Malmström et al. (2009). Obtained DNA amplicons were sequenced twice on a 454 GS FLX platform (Roche, Life Sciences) and synthetic clones underwent statistical analysis in order to identify authentic aDNA sequences for each of the individuals (Helgason et al., 2009, 2007; Malmström et al., 2012). The analyses were performed using PhyloNet v 1.0.0. software (Helgason personal communication, see Malmström et al., 2012).

Haplotype diversity was calculated according to Nei (HD) (Nei, 1987), while within population gene diversity, as well as minimum and maximum match probabilities were estimated as in Brinkmann et al. (1999). In short, the measure was calculated in order to assess the probability of obtaining different haplotypes while sampling two individuals from the same population ( $dw_{\min}$ ); the maximum ( $mw_{\max}$ ) and minimum ( $mw_{\min}$ ) probabilities of obtaining within population match.

## 3. Results

### 3.1. Carbon and nitrogen isotope analyses

All samples except the whale specimen provided sufficient collagen yields (1.1–12.0%) for carbon and nitrogen isotope measurement (Tables 2 and 3). All samples fall within the accepted C/N ratio, and carbon and nitrogen concentrations for samples of no diagenetic alteration (DeNiro, 1985; Ambrose, 1990).

The  $\delta^{13}\text{C}$  values for humans range from  $-19.5\text{‰}$  to  $-15.3\text{‰}$  ( $-16.8 \pm 0.9\text{‰}$ , mean  $\pm$  s.d.), and the  $\delta^{15}\text{N}$  values range from  $10.8$  to  $17.7\text{‰}$  ( $15.7 \pm 1.6\text{‰}$ , mean  $\pm$  s.d.) and clearly indicate a strong differentiation between individuals (Table 3). This range is significantly larger than what would be expected from a population consuming a homogeneous diet (Lovell et al., 1986), and identifies different sources of protein intake.


For individuals 5863A, 5864A and 5865A, all buried intact in double/triple burials, samples from mandibles served to control the osteologist's match between crania and postcranial remains (Table 3). All mandible samples have higher  $\delta^{15}\text{N}$  values than those

**Table 1**

MtDNA HVR1 sequence matching probabilities within ancient Flakstad, Iceland, Denmark, England and modern Norway. Comparative data was collected from the following publications: Denmark (Dissing et al., 2007; Melchior et al., 2008; Rudbeck et al., 2005), Iceland (Helgason et al., 2009), England (Töpfer et al., 2006) and Norway (Helgason et al., 2001; Krzewińska unpublished manuscript; Opdal et al., 1998; Passarino et al., 2002).

	Flakstad	Iceland	Denmark	England	Modern Norway
N	9	73	21	48	1122
HD	0.92	0.95	0.98	0.99	0.94
$dw_{\min}$	0.81	0.93	0.93	0.97	0.94
$mw_{\max}$	0.18	0.07	0.07	0.03	0.06
$mw_{\min}$	0.08	0.05	0.02	0.01	0.06

A.No	Sex	Age	Sample	% Coll.	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	%C	%N	C/N	MtDNA motif	Est. Hg
5858	F	20–25	M3 <i>Rib</i>	3.2 8.0	−16.6 −15.4	16.7 17.7	43.0 44.3	14.5 15.5	3.4 3.3	16162G	H1a
5860	F	40–45	M2 <i>L. Bone</i>	6.4 1.1	−17.4 −16.9	14.3 15.9	44.1 43.3	15.2 15.0	3.4 3.4	ND	ND
5861	M	35–45	M2 <i>L. Bone</i>	5.7 9.1	−16.3 −16.5	16.1 16.3	44.6 44.5	16.1 15.2	3.2 3.4	rCRS	H*
5863A	F	40–45	M3 Mandible	6.8 3.5	−19.2 −18.5	10.8 12.8	45.5 39.4	15.6 12.9	3.4 3.6	16069T 16126C	J*
5863B*	M	20–25	<i>L. Bone</i> <i>Femur</i>	3.7 7.8	−19.5 −15.3	12.4 17.6	41.6 44.8	15.4 15.6	3.1 3.3	16184T 16153A 16189C 16304C	H*
5864A	M	45–50	M2 Mandible	3.9 5.0	−18.9 −18.9	12.9 13.7	43.1 39.3	14.7 14.5	3.4 3.2	rCRS	H*
5864B*	M	20–35	<i>Femur sin</i> <i>Femur sin</i>	3.8 4.6	−18.9 −15.4	13.3 17.7	44.9 44.1	15.5 15.3	3.4 3.4	rCRS	H*
5864C*	?	?	<i>Femur</i>	4.0	−16.2	16.7	44.8	15.6	3.3	16134T 16240G 16286A 16356C	U4
5865A	M	25–30	M2 Mandible <i>Tibia</i>	7.1 6.9 9.7	−18.7 −16.8 −17.3	12.8 15.4 14.5	44.3 40.3 42.6	15.5 14.5 14.9	3.3 3.2 3.3	16069T 16126C 16145A 16172C 16192T 16222T 16261T	J1b1
5865B*	F	25–30?	<i>Femur dx</i>	8.7	−17.1	14.7	44.3	15.5	3.3	16192T 16256T 16270T 16291T 16325C	U5a




**Fig. 2.**  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values from teeth and bone for each individual at Flakstad. Individuals marked with "B" and "C" are all persons buried without a head in a multiple burial.

and  $-14.3\text{‰}$  in  $\delta^{13}\text{C}$ , and 15.6 and 18.1‰ in  $\delta^{15}\text{N}$ , indicating a heavy reliance on marine resources. The most elevated  $\delta^{15}\text{N}$  value is from a tooth, and could indicate that the pup was suckled during formation of the tooth. Although the  $\delta^{13}\text{C}$  value from bone is largely

comparable to that of Individuals 5858, 5863B and 5864B, the  $\delta^{15}\text{N}$  value is considerably lower, 15.6‰ as compared to 17.6–17.7‰ for the humans. This suggests that the dog was fed marine foodstuffs of lower trophic level than those consumed by people.

The mean  $\pm 2\text{s.d.}$  of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values for medieval cod from Arctic Norway (Barrett et al., 2008) were used to calculate a predicted isotopic range for someone on a diet composed exclusively of cod, including a trophic-level offset of  $+1\text{‰}$  for  $\delta^{13}\text{C}$  and  $3\text{--}5\text{‰}$  for  $\delta^{15}\text{N}$  (Fig. 3). The cod data most likely include both coastal and migratory cod – food sources which must have been available to people living at Flakstad. Six samples, from five human individuals and one dog, fall within the predicted range, suggesting a heavy reliance on marine resources. The human samples derive from either single burials, or from the decapitated individuals in multiple burials. A correspondingly high input of marine foodstuffs is also recorded in stable isotope investigations of other Norse societies (Arneborg et al., 1999; Barrett et al., 2000a,b; Barrett and Richards, 2004; Linderholm et al., 2008) and is further supported by the increasing amounts of archaeological remains from marine species in the Viking Age and following period (Barrett et al., 1999; Milner et al., 2007; Perdikaris, 1999).

Individuals buried intact in multiple burials were compared with the remaining individuals using a two-tailed *t*-test with unequal variance. A statistically significant difference ( $P \leq 0.05$ ) was shown between the two groups, both in  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values ( $P = 0.046$  and  $P = 0.011$  respectively). Only bone samples were used in calculation to ensure that each individual was represented. A comparison between individuals without heads in multiple burials and those buried in single burials showed no statistically significant difference.



**Fig. 3.** All  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values for humans and animals from Flakstad, as well as mean faunal data with error bars representing  $\pm 2\text{s.d.}$  (data from this study, North Atlantic cod presented in Barrett et al., 2008:856, pig and reindeer from Norway presented in Rosvold et al., 2010:1115). The predicted isotope range for consumers of cod from Arctic Norway are marked by the shaded square ( $+1\text{‰}$  for  $\delta^{13}\text{C}$  and  $+3\text{--}5\text{‰}$  for  $\delta^{15}\text{N}$  relative to the food source). Values from humans buried intact in multiple burials are marked with a red cross. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



### 3.2. Mitochondrial DNA

Ancient DNA analysis was undertaken in order to investigate probable maternal kinship between the individuals, as well as to establish genetic origins of people. Reproducible mitochondrial DNA sequences were obtained for 9 out of 10 individuals. Sequences from individuals 5858 and 5861 should be treated with caution as *P*-values from *c*-statistic analyses (Helgason et al., 2007) were too high to confirm ‘authenticity’. However, the low number of divergent DNA sequences among amplicons, in combination with good bone preservation, could result in less distinct deamination patterns and reduced *c*-statistic values. Therefore, we include the sequences from individuals 5858 and 5861 in our discussion.

Only three major European mtDNA haplogroups (lineages) were detected among the individuals: Hg H\*, Hg J\* and Hg U\*, reflecting relatively small sample size. All three haplogroups are of European descent (Torroni et al., 1996) and fall into previously reported Norwegian mitochondrial DNA variation (Dupuy and Olaisen, 1996; Helgason et al., 2001; Opdal et al., 1998; Passarino et al., 2002).

Three individuals (5861, 5864A and 5864B; 33%), belonging to haplogroup H\*, were carriers of DNA sequences identical to rCRS (revised Cambridge Reference Sequence) (Anderson et al., 1981; Andrews et al., 1999). The prevalence of the haplotype in present-day Europeans was estimated based on mtDNA fragment between positions 16,123–16,329 from 229 individuals from Denmark (Mikkelsen et al., 2010; Richards et al., 1996), 960 individuals from Iceland (Helgason et al., 2003, 2000; Richards et al., 1996), 234 individuals from England (Helgason et al., 2001; Richards et al., 1996) and 1122 individuals from Norway (Helgason et al., 2001; Krzewińska unpublished manuscript; Opdal et al., 1998; Passarino et al., 2002) and was found to be around 20%. In a small population like the one from Flakstad, one would, therefore, expect to identify two carriers of the sequence. The presence of three carriers inflates the observed frequency of the mtDNA motif and probably results from a stochastic sampling effect. Since all remaining individuals were carriers of different lineages, we suggest that the examined sample does not represent a maternally closely related population. In order to further investigate genetic diversity among the examined individuals, we have calculated several diversity measures as in Brinkmann et al. (1999). The probability of obtaining different haplotypes, while sampling two individuals from the same population, was calculated as a gene diversity (Nei, 1987), and was equal to 81.5% ( $dw_{min}$ ) (four in five) in the Flakstad sample. Therefore, the maximum matching probability (probability of obtaining the same haplotype while randomly sampling two individuals from Flakstad) was equal to 18.5% ( $mw_{max}$ ). The lower bounds of this estimation was 8% ( $mw_{min}$ ). Both estimates were higher than among ancient populations from Denmark, England and Iceland, as well as among modern Norwegians (Table 1). Match probabilities are influenced both by geographical range and population size. Therefore, reported comparisons between nine individuals from Flakstad and larger populations representing wide geographical locations can only serve as an approximation. Nevertheless, the analysis revealed that the lower bound for finding within population haplotype match in Flakstad (8%) was most similar to observations from most closely related populations from ancient Iceland (5%) and modern Norway (6%). Seeing that individuals from a small localized population, such as Flakstad, would be expected to share some kinship, the observed values, even if approximated, suggest the examined individuals do not represent a particularly maternally inbred population.

### 4. Discussion and conclusions

The poor contextual information available at Flakstad was supplemented by the bioarchaeological analyses, strengthening both interpretation of multiple burials and differential treatment of individuals in multiple burials as intentional and not just an effect of taphonomic processes. Stable isotope analyses were used to argue that crania and postcranial bones were correctly matched, an inference that rendered possible further investigation of diet diversity. The probable decapitation and removal of heads indicated the presence of people from different social strata – something which is further emphasized by differences in stable isotope signatures of individuals buried intact compared to those lacking crania.

With the exception of burial no 5864, ancient DNA results support this interpretation, suggesting maternal relations between individuals buried together to be unlikely. The low haplotype diversity and relatively high probability of finding matching haplotypes in the sample from Flakstad (Table 1) cannot exclude possible maternal kinship between three individuals sharing the rCRS sequence. However, the shared haplotype is the most common European lineage and minimum match probabilities in Flakstad resembled values observed in larger, closely related ancient and modern populations. Therefore, we suggest that the rCRS carriers from Flakstad were probably not maternally related, and that the seemingly high prevalence of the rCRS haplotype reflects a sampling effect in the small sample. In one case where two rCRS carriers were buried together (5864), isotopic values revealed significantly different diets for the two individuals suggesting different social standing. Social inequality implies that these individuals did not have close family relations, or at least did not share meals. Ancient DNA results presented here can only reflect direct matrilineal kinship and cannot be used to exclude other numerous kinship relations between individuals. However, in light of isotope data suggesting significantly different dietary life histories in individuals buried together, it seems unlikely the individuals were closely related.

The Flakstad burials offer an unusual insight into the way of living in a hierarchical society. It is worth noting that isotopic values of headless individuals do not differ from the values of those buried in single burials. In a society where most of the daily activities were dedicated to the acquirement and preparation of food, where food shortage and harsh winters are assumed to have been a constant threat (Sigurdsson, 2008), it would seem likely that a different diet should be detectable in people of low social standing compared to the common population. However, isotopic data in this study show quite the contrary. Despite indications that the headless people in multiple graves might represent low-status members of the population, their diet was equivalent to those buried in single burials. The persons in single burials were all buried in a seemingly respectful way and accompanied by grave gifts, and are interpreted as representatives of the free population. They all had a diet combining marine and terrestrial sources, and a significant part of their diet consisted of marine foodstuffs – as did that of the headless individuals. It is of course possible that there might have been other dietary differences between social groups, through preparation and possibly the amount of protein, rather than different food sources. The complete individuals from multiple burials, however, stand out as a distinct group and may be perceived as persons of a special social status, emphasized by their distinctly different diet already evident during childhood. The lack of high status artifacts in multiple burials – although uncertain due to agricultural disturbance – could indicate that the complete individuals were not necessarily wealthy, but special in another sense. An alternative explanation might be previous disturbance,

not just as a result of agriculture, but also the possible plunder and removal of precious objects. Either way, the individuals buried intact seem to have belonged to separate social strata, treated differently than others in death as well as in life. Due to the poor contextual information from the site these observations could not have been made without information hidden in the human remains.

To conclude, the multidisciplinary approach to the skeletal remains from Flakstad has revealed some intriguing patterns. Results from stable isotope analyses show that individuals in multiple burials most likely were intentionally placed in the same burial, given the pattern in which the only person buried intact in each burial, had distinct isotope values. Thus, persons sharing a grave had distinctly different diets during their lifetime and were unlikely to share maternal kinship. A reasonable explanation for these observations could be that persons buried headless may have been slaves accompanying their masters in the grave. This interpretation corresponds well with other double burials from the Norse World with similar features, where decapitated and sometimes headless people were deposited as grave gifts. The resemblance in diet between headless persons and individuals buried in single burials was unexpected and calls for further investigation in the future. The present study indicates that also other double burials should be investigated using a bioarchaeological approach. This material offers an exceptional opportunity to study the subpopulation groups of the Norse, and their way of living in a divided and hierarchical world.

## Acknowledgements

We wish to express our gratitude to Jan Bill, head of the SIGMA project, who made this study possible, Lotte Hedeager, Marcin Wojewodzic and Gro Bjørnstad for valuable reading and feedback on the manuscript, Per Holck for providing assistance and advice with the human remains, Agnar Helgason for making the PhyloNet program available to us, Anne Karin Hufthammer for assisting with animal reference material, Elin T. Brødholt for commenting on osteological specifications in Table 3, Heike Siegmund at the Stable Isotope Laboratory (SIL), for running the EA-IRMS, Knut Helskog and Aud Ahlquist at Tromsø University Museum for permission and assistance and The National Committee for Research Ethics on Human Remains for evaluation of the project. We also want to thank two anonymous referees for thorough and valuable feedback on the manuscript.

## References

- Ambrose, S.H., 1990. Preparation and characterization of bone and tooth collagen for isotopic analysis. *J. Archaeol. Sci.* 17, 431–451.
- Ambrose, S.H., Norr, L., 1993. Experimental evidence for the relationship of carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: Lambert, J.B., Grupe, G. (Eds.), *Prehistoric Human Bone: Archaeology at the Molecular Level*. Springer-Verlag, Berlin, pp. 1–37.
- Ambrose, S.H., Butler, B.M., Hanson, D.B., Hunter-Anderson, R.L., Krueger, H.W., 1997. Stable isotope analysis of human diet in the Marianas archipelago, Western Pacific. *Am. J. Phys. Anthropol.* 104, 343–361.
- Andersen, H., 1960. Hovedstaden i Riget. Nationalmuseets arbejdsmark, 1960. Nationalmuseet, København.
- Anderson, S., Bankier, A.T., Barrell, B.G., de Bruijn, M.H., Coulson, A.R., Drouin, J., Eperon, I.C., Nierlich, D.P., Roe, B.A., Sanger, F., Schreier, P.H., Smith, A.J., Staden, R., Young, I.G., 1981. Sequence and organization of the human mitochondrial genome. *Nature* 290, 457–465.
- Andrews, R.M., Kubacka, I., Chinnery, P.F., Lightowlers, R.N., Turnbull, D.M., Howell, N., 1999. Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nat. Genet.* 23, 147.
- Arneborg, J., Heinemeier, J., Lynnerup, N., Nilsen, H.L., Rud, N., Sveinbjörnsdóttir, A.E., 1999. Change of diet of the Greenland Vikings determined from stable isotope analysis and  $^{14}\text{C}$  dating of their bones. *Radiocarbon* 41 (2), 157–168.
- Bandelt, H.J., Macaulay, V., Richards, M. (Eds.), 2006. *Human Mitochondrial DNA and the Evolution of Homo Sapiens*. Springer-Verlag, Berlin, Heidelberg.
- Barrett, J.H., Beukens, R., Simpson, I., Ashmore, P., Poaps, S., Huntley, J., 2000a. What was the Viking age and when did it happen? A view from Orkney. *Nor. Archaeol. Rev.* 33.
- Barrett, J.H., Nicholson, R.A., Cerón-Carrasco, R., 1999. Archaeo-ichthyological evidence for long-term socioeconomic trends in Northern Scotland: 3500 BC to AD 1500. *J. Archaeol. Sci.* 26, 353–388.
- Barrett, J.H., Beukens, R.P., Brothwell, D.R., 2000b. Radiocarbon dating and marine reservoir correction of Viking Age Christian burials from Orkney. *Antiquity* 74, 537–543.
- Barrett, J.H., Richards, M.P., 2004. Identity, gender, religion and economy: new isotope and radiocarbon evidence for marine resource intensification in early historic Orkney, Scotland, UK. *Eur. J. Archaeol.* 7 (3), 249–271.
- Barrett, J.H., Johnstone, C., Harland, J., Neer, W.V., Erynck, A., Makowiecki, D., Heinrich, D., Hufthammer, A.K., Enghoff, I.B., Amundsen, C., Christiansen, Jones, A.K.G., Locker, Hamilton-Dyer, S., Jonsson, Leif, Løugas, L., Roberts, C., Richards, M., 2008. Detecting the medieval cod trade: a new method and first results. *J. Archaeol. Sci.* 35, 850–861.
- Bocherens, H., Drucker, D., 2003. Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems. *Int. J. Osteoarchaeol.* 13, 46–53.
- Brinkmann, C., Forster, P., Schürenkamp, M., Horst, J., Rolf, B., Brinkmann, B., 1999. Human Y-chromosomal STR haplotypes in a Kurdish population sample. *Int. J. Leg. Med.* 112, 181–183.
- Brown, T.A., Nelson, D.E., Vogel, J.S., Southon, J.R., 1988. Improved collagen extraction by modified longin method. *Radiocarbon* 30 (2), 171–177.
- Chisholm, B.S., Nelson, D.E., Schwarcz, H.P., 1982. Stable-carbon isotope ratios as a measure of marine versus terrestrial protein in ancient diets. *Sci. New Ser.* 216 (4550), 1131–1132.
- DeNiro, M.J., 1985. Postmortem preservation and alteration of in vivo bone collagen isotope ratios in relation to paleodietary reconstruction. *Nature* 317, 806–809.
- Dissing, J., Binladen, J., Hansen, A., Sejrsen, B., Willerslev, E., Lynnerup, N., 2007. The last Viking King: a royal maternity case solved by ancient DNA analysis. *Forensic Sci. Int.* 166, 21–27.
- Dupuy, B.M., Olaisen, B., 1996. mtDNA sequences in Norwegian Saami and main populations. In: Carracedo, A., Brinkmann, B., Bär, W. (Eds.), *Advances in Forensic Haemogenetics*, first ed. Springer Verlag, Berlin, Heidelberg, pp. 23–25.
- Eriksson, G., Lidén, K., 2013. Dietary life histories in Stone age northern Europe. *J. Anthropol. Archaeol.* 32 (3), 288–302.
- Hagelberg, E., Clegg, J.B., 1991. Isolation and characterization of DNA from archaeological bone. *Proc. R. Soc. Lond. B.* 244, 45–50.
- Hedges, R., Clement, J.G., Thomas, D.L., O'Connell, T.C., 2007. Collagen turnover in the adult femoral mid-shaft: modeled from anthropogenic radiocarbon tracer measurements. *Am. J. Phys. Anthropol.* 133, 808–816.
- Helgason, A., Hickey, E., Goodacre, S., Bosnes, V., Stefánsson, K., Ward, R., Sykes, B., 2001. mtDNA and the islands of the North Atlantic: estimating the proportions of Norse and Gaelic ancestry. *Am. J. Hum. Genet.* 68, 723–737.
- Helgason, A., Nicholson, G., Stefánsson, K., Donnelly, P., 2003. A reassessment of genetic diversity in Icelanders: strong evidence from multiple loci for relative homogeneity caused by genetic drift. *Ann. Hum. Genet.* 67, 281–297.
- Helgason, A., Lalueza-Fox, C., Ghosh, S., Sigurðardóttir, S., Sampietro, M.L., Gigli, E., Baker, A., Bertranpetit, J., Árnadóttir, L., Thorsteinsdóttir, U., Stefánsson, K., 2009. Sequences from first settlers reveal rapid evolution in Icelandic mtDNA pool. *PLoS Genet.* 5, e1000343.
- Helgason, A., Pálsson, S., Lalueza-Fox, C., Ghosh, S., Sigurðardóttir, S., Baker, A., Hrafnkelsson, B., Árnadóttir, L., Thorsteinsdóttir, U., Stefánsson, K., 2007. A statistical approach to identify ancient template DNA. *J. Mol. Evol.* 65, 92–102.
- Helgason, A., Sigurðardóttir, S., Gulcher, J.R., Ward, R., Stefánsson, K., 2000. mtDNA and the origin of the Icelanders: deciphering signals of recent population history. *Am. J. Hum. Genet.* 66, 999–1016.
- Hemmendorff, O., 1984. Människooffer. Ett innslag i jernalderens gravritualer, belyst av ett fynd i Bollstanäs, Uppland. *Fornvännen* 79, 4–12.
- Holck, P., 1997. Cremated Bones. A Medical-anthropological Study of an Archaeological Material on Cremation Burials. Anatomical Institute, University of Oslo.
- Holck, P., 1983. Rapport for skjelettmateriale fra Flakstad. Bergen Museum/Anatomical Institute, University of Oslo (Unpublished).
- Holmquist Olausson, L., 1990. "Älgmannen" från Birka: Presentation av en nyligen undersökt krigargrav med människooffer. *Fornvännen* 85, 175–182.
- Jones, J.R., Mulville, J.A., McGill, R.A.R., Evershed, R.P., 2012. Paleoenvironmental modelling of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in the North Atlantic Islands: understanding past marine resource use. *Rapid Commun. Mass. Spectrom.* 26, 2399–2406.
- Linderholm, A., Hedenstierna Jonson, C., Svensk, O., Lidén, K., 2008. Diet and status in Birka: stable isotopes and grave goods compared. *Antiquity* 82, 446–461.
- Lovell, N.C., Nelson, D.E., Schwarcz, H.P., 1986. Carbon isotope ratios in paleodiet: lack of age or sex effect. *Archaeometry* 28 (1), 51–55.
- Malmström, H., Gilbert, M.T.P., Thomas, M.G., Brandström, M., Storå, J., Molnar, P., Andersen, P.K., Bendixen, C., Holmlund, G., Götherström, A., Willerslev, E., 2009. Ancient DNA reveals lack of continuity between Neolithic hunter-gatherers and contemporary Scandinavians. *Curr. Biol.* 19, 1758–1762.
- Malmström, H., Vretemark, M., Tillmar, A., Durling, M.B., Skoglund, P., Gilbert, M.T.P., Willerslev, E., Holmlund, G., Götherström, A., 2012. Finding the founder of Stockholm – a kinship study based on Y-chromosomal, autosomal and mitochondrial DNA. *Ann. Anat.* 194, 138–145.
- Melchior, L., Kivisild, T., Lynnerup, N., Dissing, J., 2008. Evidence of authentic DNA from Danish Viking Age skeletons untouched by humans for 1,000 years. *PLoS One*, e2214.

- Mikkelsen, M., Sørensen, E., Rasmussen, E.M., Morling, N., 2010. Mitochondrial DNA HV1 and HV2 variation in Danes. *Forensic Sci. Int. Genet.* 4, e87–88.
- Milner, N., Barrett, J.H., Welsh, J., 2007. Marine resource intensification in Viking age Europe: the Molluscan evidence from Quoygrew, Orkney. *J. Archaeol. Sci.* 34 (9), 1461–1472.
- Minagawa, M., Wada, E., 1984. Stepwise enrichment of  $^{15}\text{N}$  along food chains: further evidence and the relation between  $\delta^{15}\text{N}$  and animal age. *Geochim. Cosmochim. Acta* 48, 1135–1140.
- Nei, M., 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- Opdal, S.H., Rognum, T.O., Vege, Å., Stave, A.K., Dupuy, B.M., Egeland, T., 1998. Increased number of substitutions in the D-loop of mitochondrial DNA in the sudden infant death syndrome. *Acta Paediatr.* 87, 1039–1044.
- Passarino, G., Cavalleri, G.L., Lin, A.A., Cavalli-Sforza, L.L., Børresen-Dale, A.L., Underhill, P.A., 2002. Different genetic components in the Norwegian population revealed by the analysis of mtDNA and Y chromosome polymorphisms. *Eur. J. Hum. Genet.* 10, 521–529.
- Perdikaris, S., 1999. From Chiefly provisioning to commercial fishery: long-term economic change in Arctic Norway. *World Archaeol.* 30 (3), 388–402.
- Pääbo, S., 1989. Ancient DNA: extraction, characterization, molecular cloning, and enzymatic amplification. *Proc. Natl. Acad. Sci. U.S.A* 86, 1939–1943.
- Ramskou, T., 1965. *Vikingerne ofrede mennesker*. Nationalmuseets Arbejdsmark 1963–1965. Nationalmuseet, København.
- Richards, M.B., Sykes, B.C., Hedges, R.E.M., 1995. Authenticating DNA extracted from ancient skeletal remains. *J. Archaeol. Sci.* 22, 291–299.
- Richards, M., Corte-Real, H., Forster, P., Macaulay, V., Wilkinson-Herbots, H., Demaine, A., Papiha, S., Hedges, R., Bandelt, H.J., Sykes, B., 1996. Paleolithic and Neolithic lineages in the European mitochondrial gene pool. *Am. J. Hum. Genet.* 59, 185–203.
- Rosvold, J., Halley, D.J., Hufthammer, A.K., Minagawa, M., Andersen, R., 2010. The rise and fall of wild boar in a northern environment: evidence from stable isotopes and subfossil finds. *Holocene*, 1113–1121.
- Rudbeck, L., Gilbert, M.T.P., Willerslev, E., Hansen, A.J., Lynnerup, N., Christensen, T., Dissing, J., 2005. mtDNA analysis of human remains from an early Danish Christian cemetery. *Am. J. Phys. Anthropol.* 128, 424–429.
- Sandmo, A.K., 1985a. Flakstad Prestegård -Innberetninger/rapporter. In: *Utgravninger v/Anne Karine Sandmo 1980–85*. Tromsø Museum, Tromsø.
- Sandmo, A.K., 1985b. Gravfelt og bosetningsområder fra eldre og yngre jernalder på Flakstad i Lofoten. Foreløpige gravingsresultater og noen hypoteser. In: Engelstad, E., Holm-Olsen, I.M. (Eds.), *Arkeologisk Feltarbeid i Nord-Norge 1984*. University of Tromsø. Tromsø, Kulturhistorie no. 5.
- Schoeninger, M.J., DeNiro, M.J., Tauber, H., 1983. Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Sci. New Ser.* 220 (4604), 1381–1383.
- Sealy, J., Armstrong, R., Schrire, C., 1995. Beyond lifetime averages: tracing life histories through isotopic analysis of different calcified tissues from archaeological human skeletons. *Antiquity* 69, 290–300.
- Sigurdsson, J.V., 2008. *Det Norrøne Samfunnet: Vikingen, Kongen, Erkebiskopen Og Bonden*. Pax Forlag, Oslo.
- Skre, D., 1998. *Herredømmet. Bosetning og besittelse på Romerrike 200–1350 e. Kr.* Scandinavian University Press, University of Oslo.
- The Schreiner Collection Database, University of Oslo.** (accessed 23.08.13.).
- Torroni, A., Huoponen, K., Francalacci, P., Petrozzi, M., Morelli, L., Scozzari, R., Obinu, D., Savontaus, M.L., Wallace, D.C., 1996. Classification of European mtDNAs from an analysis of three European populations. *Genetics* 144, 1835–1850.
- Töpf, A.L., Gilbert, M.T., Dumbacher, J.P., Hoelzel, A.R., 2006. Tracing the phylogeography of human populations in Britain based on 4th–11th century mtDNA genotypes. *Mol. Biol. Evol.* 23, 152–161.
- Zachrisson, T., 2003. De osynliggjorda. Om trälar i arkeologisk forskning. In: Lindquist, T., Myrdal, J. (Eds.), *Trälar. Ofria i agrarsamhället från vikingatid till medeltid*. Nordiska museets förlag, Stockholm.